PATIENT INFORMATION LEAFLET

HOW TO USE YOUR STEROID CREAMS

for eczema flares (f) and maintenance (m)

WHEN YOU HAVE A FLARE

1. SPEAK TO YOUR GP for a flare (f) prescription:

They will give you a strong steroid.

Use this:

Twice daily for 7 days

then, Once daily for 7 days

then, Twice a week for 14 days

Is this eczema?

red itchy weepy

relapses and remits

havfever/ asthma run in the family

Face

2.5 FTU

chest

2.5 FTU

stomach

2.5 FTU groin

F

AFTU

THE FACE

Steroid-free option for treating facial eczema (discuss with your GP)

> 2.5 Fingertip Units Tacrolimus cream

> > Use this:

Twice daily for flares (f) Twice a week for maintenance (m)

2. SPEAK TO YOUR GP for a maintenance (m) prescription:

They will give you a weak steroid.

Use this:

Twice a week ongoing

2.5 FTU

FINGERTIP UNITS PER APPLICATION

Use the Fingertip Unit (FTU) rule

1 FTU = amount of cream/ointment squeezed from the tip of an adult finger to the first crease.

Example doses:

Scalp: 1.5 FTU Face and neck: 2.5 FTU One arm: 2.5 FTU One hand: 1.5 FTU Leg and foot: 8 FTU

For children, use half of the above

EMOLIENTS

Apply emollients after washing, whilst the skin is still damp, and 60 minutes before the steroid

QV cream - morning creams are less effective, but feel lighter on the skin

Dermol 500 - to wash has antimicrobial properties

Diprobase ointment - evening ointments are effective for dry skin but are greasy

EMOLIENTS ARE FLAMMABLE, DO NOT USE NEAR FIRE

WARNING

Do not suddenly stop using your steroid. Otherwise, you may experience a rebound flare of symptoms.

Long-term steroid use can lead to skin thinning.